



**RHODE ISLAND'S RECOMMENDED CHILDHOOD IMMUNIZATION SCHEDULE FOR ROUTINE<sup>1</sup>  
VACCINATION USING STATE-SUPPLIED VACCINES<sup>2</sup>**

VACCINE	MONTHS							YEARS											
	Birth	2	4	6	12	15	18	19-23	4-6	11-12	16	18							
<b>Hepatitis B</b>	<b>Hep B</b>	<i>Pediarix</i> (DTaP-Hep B-IPV) 1 dose at 2, 4, & 6 months																	
<b>DTaP/Tdap</b> (Diphtheria, Tetanus, Pertussis)													<i>Infanrix</i> DTaP			<i>Kinrix</i> (DTaP-IPV)	<i>Boostrix</i> Tdap		
<b>IPV</b> (Polio)																			
<b>Hib</b> (Haemophilus influenzae type B)		<i>PedvaxHIB</i> Hib	<i>PedvaxHIB</i> Hib		<i>PedvaxHIB</i> Hib														
<b>PCV13</b> (Pneumococcal conjugate)		<i>Prevnar</i> PCV13	<i>Prevnar</i> PCV13	<i>Prevnar</i> PCV13	<i>Prevnar</i> PCV13														
<b>RV</b> (Rotavirus)		<i>Rotarix</i> RV	<i>Rotarix</i> RV																
<b>MMR</b> (Measles, Mumps, Rubella)					MMR				MMRV										
<b>Chickenpox</b> (Varicella)					<i>Varivax</i> Varicella														
<b>Hepatitis A</b>					Hep A 2 doses (6 months apart)														
<b>MCV4</b> (Meningococcal conjugate)										<i>Menactra</i> MCV4	<i>Menactra</i> MCV4								
<b>MenB</b> (Meningococcal serotype B)											<i>MenB</i> 2 doses at least 1 month apart								
<b>HPV9<sup>3</sup></b> (Human papillomavirus)										<i>Gardasil 9</i> : 2 doses (0, 6-12 months) at 11-12 yrs; age 15 & older 3 dose series (0, 1-2, 6 months)									
<b>Influenza</b>				Influenza (1 or 2 doses as indicated) Yearly during flu season															

<sup>1</sup> Refer to CDC's *Recommended Childhood and Adolescent Immunization Schedule* for vaccinating high risk children at: <http://www.cdc.gov/vaccines/schedules/hcp/imz/child-adolescent.html>

<sup>2</sup> This schedule lists state-supplied childhood vaccines and is consistent with the recommended age range of CDC's *Childhood and Adolescent Immunization Schedule* (<http://www.cdc.gov/vaccines/schedules/hcp/child-adolescent.html>). CDC recommendations for individual vaccines are available at [www.cdc.gov/vaccines/recs/default.htm](http://www.cdc.gov/vaccines/recs/default.htm).

<sup>3</sup> State-supplied 9vHPV vaccine is recommended for routine use in males and females beginning at 11-12 years with catch-up vaccination through 18 years of age.